

CLAIMS

- Sub B1
- 1 1. A method of preventing thrombosis formation on a liquid-containing surface of a liquid
2 delivery system, the liquid delivery system being connected to a patient for delivery of a
3 liquid to said patient, the method comprising a regimen selected from the group consisting of:
4 A) contacting said surface with a thrombosis-preventing liquid containing
5 taurolidine, taurultam or a mixture thereof, said thrombosis-preventing liquid
6 further containing an anticoagulant agent; and
7 B) first contacting said surface with a solution containing a thrombosis-
8 preventing amount of an anticoagulant agent, and thereafter contacting said
9 surface with a solution containing taurolidine, taurultam or a mixture thereof.

- Sub C2
- 1 2. The method of claim 1 wherein the solution or liquid containing taurolidine, taurultam or
2 mixture thereof is contacted with said surface for at least about 1 hour.
3 3. The method of claim 2 wherein said solution or liquid containing taurolidine, taurultam or
4 mixture thereof is sealed in said delivery system for a period of at least 12 hours.
5 4. The method of claim 3 wherein said solution or liquid containing taurolidine, taurultam or
6 mixture thereof which is sealed in said delivery system, is replaced at least about daily.
7 5. The method of claim 1 wherein, when first contacting said surface with the anticoagulant
8 solution, said surface is flushed with said anticoagulant-containing solution.
9 6. The method of claim 5 wherein the solution containing taurolidine, taurultam or a mixture
thereof is contacted with said surface for at least about 1 hour.
1 7. The method of claim 6 wherein said solution containing taurolidine, taurultam or a
2 mixture thereof is sealed in said delivery system for a period of at least about 12 hours.
3 8. The method of claim 7 wherein the solution containing taurolidine, taurultam or a mixture
4 thereof which is sealed in said delivery system is replaced at least about daily.

13

1 17. The composition of claim 16 wherein said anticoagulant agent is selected from the group
2 consisting of aprotinin, hirudin, desirudin, danaparoid, danaparoid-sodium, pentosan,
3 pentosanpolysulfate-sodium, ticlopidine, clopidogrel, and mixtures thereof.

1 18. The composition of claim 16 wherein said anticoagulant agent is heparin.

1 19. The composition of claim 16 wherein said anticoagulant is sodium citrate.

1 20. The composition of claim 16 wherein said turoolidine is present in an amount of from
2 about 0.5 to about 3% by weight or said taurultam is present in an amount of from about 1 to
3 about 7.5% by weight, and said anticoagulant agent is present in an amount within a range of
4 from about 0.01 to about 5% by weight.

add
B2

009100-85522560